

HVLab Multi-channel Plethysmograph:

Lewis CH (1996) The development of a multi-channel plethysmograph, Proceedings from the United Kingdom Informal Group Meeting on Human Responses to Vibration, MIRA, 18-20 September, 1-12.

Lindsell CJ, Griffin MJ (1998) Standardised diagnostic methods for assessing components of the hand-Arm vibration syndrome. Published by HSE Books, ISBN 0-7176-1640-1.

Lindsell CJ, Griffin MJ (1999) Thermal thresholds, vibrotactile thresholds and finger systolic blood pressures in dockyard workers exposed to hand-transmitted vibration. *Int Arch Occup Environ Health* 1999, 72: 377-386

Lindsell CJ, Griffin MJ (2000) A standardised test battery for assessing vascular and neurological components of the hand-arm vibration syndrome. In, *Proceedings of the 8th International Conference on Hand-Arm Vibration, Umeå, Sweden 09 - 12 Jun 1998*. Solna, Sweden, Arbets Miljo Institutet, 133-141.

Bovenzi M, Lindsell CJ, Griffin MJ (2000) Acute vascular responses to the frequency of vibration transmitted to the hand. *Occup Environ Med* 2000, 57:422-430

Bovenzi M, Lindsell CJ, Griffin MJ (2001) Response of finger circulation to energy equivalent combinations of magnitude and duration of vibration. *Occup Environ Med* 2001, 58:185-193

Lindsell CJ, Griffin MJ (2002) Normative data for vascular and neurological tests of the hand-arm vibration syndrome. *Int Arch Occup Environ Health* 2002, 75: 43-54

Lindsell CJ (2005) Test battery of assessing vascular disturbances of fingers. *Environ Health Med* 2005, 10:341-350

Bovenzi M, Welsh AJL, Griffin MJ (2007) Effect of prior exposure to hand-transmitted vibration on cold response of digital arteries. *Int Arch Occup Environ Health* 2007, 80:281-289

Welsh AJL, Griffin MJ (2008) Normal values for finger systolic blood pressures in males and females. *Int Arch Occup Environ Health* 2008, 81:625-632

Bovenzi M, Alessandrini B, Manicini R, Cannavà MG, Centi L (1998) A prospective study of the cold response of digital vessels in forestry workers exposed to saw vibration. *Int Arch Occup Environ Health* 1998, 71:493-498

Bovenzi M (2002) Finger systolic blood pressure indices for the diagnosis of vibration-induced white finger. *Int Arch Occup Environ Health* 2002, 75: 20-28

Kurozawa Y, Nasu Y, Hosoda T, Nose T (2002) Long-term follow up study on patients with vibration-induced white finger. *Occup Environ Med* 2002, 44, 12, 1203-1206

Kurozawa Y, Nasu Y (2005) Factors influencing finger systolic blood pressure test for diagnosis of vibration-induced white finger. *Environ Health and Med* 2005, 10: 366-370

Bovenzi M, Vedova AD, Negro C (2005) A follow up study of vibration induced white finger in compensation claimants. *Occup Environ Med* 2005, 62: 237-242

- Bovenzi M (2008) A follow up study of vascular disorders in vibration-exposed forestry workers. *Int Arch Occup Environ Health* 2008, 81: 401-408
- Negro C, Rui F, D'Agostin F, Bovenzi M (2008) A longitudinal study of finger systolic blood pressure and exposure to hand-transmitted vibration. *Int Arch Occup Environ Health* 2008, 81:613-623
- Bovenzi M, D'Agostin F, Rui F, Negro C (2008) Use of color charts for the diagnosis of finger whiteness in vibration-exposed workers. *Int Arch Occup Environ Health* 2008, 81:633-638
- Fujiwara Y, Yoshino S, Nasu Y (2008) Simultaneous observation of zero-value of FSBP% and Raynaud's phenomenon during cold provocation in vibration syndrome. *J Occup Health* 2008, 50:75-78